

April 2002

The neutrino puzzle in the light of SNO

R. Foot¹ and R. R. Volkas²

*School of Physics
Research Centre for High Energy Physics
The University of Melbourne
Victoria 3010 Australia*

Abstract

SNO's neutral current measurement has added a new piece to the emerging neutrino physics puzzle. Putting together the presently available experimental information, an essentially unique picture emerges: The solar neutrino anomaly is explained by $\nu_e \rightarrow \nu_\tau$ oscillations, the atmospheric neutrino anomaly is explained by $\nu_\mu \rightarrow \nu_s$ and the LSND data are explained by $\nu_e \rightarrow \nu_\mu$. This scheme will be tested by future experiments: MiniBooNE will test the oscillation explanation of the LSND anomaly, while the long baseline experiments will discriminate between the $\nu_\mu \rightarrow \nu_s$ and $\nu_\mu \rightarrow \nu_\tau$ possibilities for resolving the atmospheric anomaly (confirming or disconfirming the Super-Kamiokande result that the latter is favoured over the former).

¹foot@physics.unimelb.edu.au

²r.volkas@physics.unimelb.edu.au

Over the last decade or so, important insight into the behaviour of neutrinos has been gained from a number of important experiments. We now have strong evidence for three neutrino anomalies:

- ν_μ disappearance: Super-Kamiokande has established that the number of up-going contained ν_μ events is approximately half that of the down-going events, while there is no discernible up-down asymmetry in the ν_e -induced signal. These data can be explained by maximal oscillation of ν_μ into another species that cannot be ν_e [1]. Results from Soudan II support this conclusion[2].
- SNO has now measured both the charged-current (CC) and neutral current (NC) solar neutrino rates, finding the flux ratio $\phi^{NC}/\phi^{CC} = 2.9 \pm 0.4$ [3]. These measurements are consistent with other solar neutrino experiments[4], and suggest large mixing angle³ oscillations of ν_e into ν_μ or ν_τ , with $\sin^2 2\theta \gtrsim 0.7$ and a range of possible δm^2 . (If all solar neutrino results are taken at face value, then $\delta m^2 \sim 10^{-4} \text{ eV}^2$ is favoured. The SNO data on their own allow a wider range.)
- The LSND collaboration has obtained strong evidence for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations via an appearance experiment. They find $\delta m^2 \sim 1 \text{ eV}^2$ and $\sin^2 2\theta \sim 10^{-2}$ [6].

If we assume that no light neutrinos exist beyond the known ν_e, ν_μ and ν_τ active flavours, then one is led to the embarrassing conclusion that at least one piece of evidence must be discarded since all three neutrino anomalies cannot be reconciled. It is usually assumed that the LSND anomaly is due to an unidentified error rather than oscillations, with the solar and atmospheric neutrino anomalies then used as evidence for a close-to bi-maximal mixing pattern amongst the three active neutrinos[7]. Unfortunately, no compelling theoretical explanation for a bi-maximal-like pattern has emerged. On the other hand, there are simple schemes, including mirror symmetry and pseudo-Dirac structure, that can explain the existence of *two*-flavour maximal mixing. Furthermore, the mirror symmetry idea actually predicts the existence of light effectively sterile neutrinos, so it is worth taking the possible existence of such additional species seriously.

It is quite possible that the three neutrino anomalies are all explained by oscillations. If this is the case, then it is interesting that we are led

³Although the small-angle MSW solution is disfavoured by the apparently undistorted nature of the ^8B neutrino spectrum, the overall χ^2_{min} of 99.9 for 78 degrees of freedom, implies a C.L. of 4.8% [5], which is acceptable, so a small-angle solution is still possible.

to an essentially unique picture: LSND implies small angle $\nu_e \rightarrow \nu_\mu$ with $\delta m^2 \sim 1 \text{ eV}^2$, SNO and the other solar neutrino experiments then suggest that ν_e oscillates into ν_τ with $\sin^2 2\theta \gtrsim 0.7$, for a range of δm^2 . This leaves atmospheric ν_μ disappearance to be explained by approximately (or indeed exactly) maximal $\nu_\mu \rightarrow \nu_s$ oscillations. We formalise these observations by proposing the following hypothesis: *The fundamental theory of neutrino mixing, whatever it is, features (i) large (or even maximal) ν_μ - ν_s mixing, (ii) small-angle active-active mixing except for the ν_e - ν_τ channel which is large.*

Interestingly, this scheme has *not* been considered previously in the literature, perhaps because the Super-Kamiokande atmospheric data prefer the $\nu_\mu \rightarrow \nu_\tau$ channel *in comparison with* the $\nu_\mu \rightarrow \nu_s$ possibility. However, what really needs to be done to examine our hypothesis is not to compare the χ^2_{min} of the $\nu_\mu \rightarrow \nu_s$ solution with the χ^2_{min} of the $\nu_\mu \rightarrow \nu_\tau$ solution, but rather, to compute the allowed C.L. of the $\nu_\mu \rightarrow \nu_s$ hypothesis and check whether the goodness-of-fit is reasonable. When this is done, the $\nu_\mu \rightarrow \nu_s$ solution *is* found to be reasonable, being only mildly disfavoured, at the $1.5 - 3\sigma$ level, depending on how the data are analysed[8, 9]. For example, a recent[10] Super-Kamiokande global fit for $\nu_\mu \rightarrow \nu_s$ oscillations has found a χ^2_{min} of 222.7 for 190 degrees of freedom. This corresponds to an allowed C.L. of 5.5%, that is, it is only mildly disfavoured at about the 2σ level.

There are two possibilities for the mass hierarchy: either ν_e, ν_τ are $\sqrt{\delta m^2_{LSND}} \sim 1 \text{ eV}$ heavier than ν_μ, ν_s or vice-versa. If the ν_e, ν_τ pair is heavier, then it might put the ν_e mass, assuming it is Majorana, in the range accessible to neutrinoless double beta decay experiments.⁴

In the nearer future, the two most important tests will come from the MiniBooNE experiment and the long baseline experiments MINOS and CNGS. MiniBooNE has been designed to check the oscillation explanation of the LSND result, and with it the indirect evidence for a light sterile flavour. Through NC or τ -appearance measurements, the up-coming long baseline experiments hope to discriminate between the $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ channels used to resolve the atmospheric anomaly. This will provide a terrestrial check of the Super-Kamiokande result that the ν_τ possibility is favoured.

On the theoretical front, note that this four-neutrino phenomenological scheme can be embedded into interesting six-neutrino theories, such as provided by the mirror matter model[11]. Recall that this theory predicts that each ordinary neutrino, ν_α where $\alpha = e, \mu, \tau$, is maximally mixed with an effectively sterile mirror partner ν'_α . The interfamily mixing pattern, the absolute mass scales, and the $\nu_\alpha - \nu'_\alpha$ mass splittings are all free parameters.

⁴However, if the large-angle between ν_e and ν_τ is due to an approximate pseudo-Dirac character, then neutrinoless double beta decay may occur too slowly to be seen.

The ordinary-mirror neutrino sector will reduce to the above four-neutrino scenario if the oscillation lengths of the $\nu_e \rightarrow \nu'_e$ and $\nu_\tau \rightarrow \nu'_\tau$ modes are greater than the Earth-Sun distance at solar neutrino energies. (It might be possible to relax this assumption, but further work needs to be done.) The importance of this possible theoretical basis for the scheme is twofold. First, the presence of a light effectively sterile neutrino (the ν'_μ) is explained, because mirror neutrinos are light for exactly the same reason that ordinary neutrinos are light (whatever that reason might be). Second, the atmospheric mixing is predicted to be exactly pairwise maximal, unlike the $\nu_e - \nu_\tau$ mixing angle which could be just accidentally large. These features can also be more stringently tested in the future. For example, the Japan Hadron Facility to Super-Kamiokande superbeam proposal cites an experimental precision of about 0.01 on the atmospheric $\sin^2 2\theta$ as achievable [12]. KAMLAND may soon provide more information on the solar mixing angle.

In summary, the most important features of the three neutrinos anomalies – solar, atmospheric and LSND – can be explained through large angle $\nu_e \rightarrow \nu_\tau$, exactly or approximately maximal $\nu_\mu \rightarrow \nu_s$, and small angle $\nu_e \rightarrow \nu_\mu$ oscillations, respectively. The new information from SNO, when combined with the LSND result, suggests that the solar channel is predominantly ν_τ . Important aspects of this scenario will be tested soon by MiniBooNE and the long baseline experiments. When embedded within the mirror matter model, an important difference between the large solar and the large atmospheric mixing angles emerges: the former is large for possibly accidental reasons, while the latter is rigorously maximal for theoretical reasons. Future precision measurements of ν_μ disappearance in superbeam experiments will test that possibility further.

References

- [1] Super-Kamiokande Collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett. 82, 1562 (1998); Phys. Lett. B436, 33 (1998); Phys. Lett. B433, 9 (1998).
- [2] Soudan 2 Collaboration, W. W. Allison *et al.*, Phys. Lett. B449, 137 (1999).
- [3] SNO Collaboration, Q. R. Ahmad *et al.*, nucl-ex/0204008; nucl-ex/0204009; Phys. Rev. Lett. 87, 071301 (2001).
- [4] Homestake Collaboration, B. T. Cleveland *et al.*, Astrophys. J. 496, 505 (1998); Kamiokande Collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett. 77, 1683 (1996); Super-Kamiokande Collaboration, Phys. Rev. Lett. 86, 5651 (2001); Sage Collaboration, J. N. Abdurashitov, *et al.*, Phys. Rev. Lett. 83, 4686 (1999); Gallex Collaboration, W. Hampel *et al.*, Phys. Lett. B447, 127 (1999); GNO Collaboration, M. Altann *et al.*, Phys. Lett. B490, 16 (2000).
- [5] See Table 1 of P. C. de Holanda and A. Yu. Smirnov, hep-ph/0205241.
- [6] LSND Collaboration, C. Athanassopoulos *et al.*, Phys. Rev. Lett. 81, 1774 (1998); Phys. Rev. C58, 2489 (1998).
- [7] V. Barger *et al.*, Phys. Lett. B437, 107 (1998); A. Baltz, A. S. Goldhaber and M. Goldhaber, Phys. Rev. Lett. 81, 5730 (1998); F. Vissani, hep-ph/9708483; D. V. Ahluwalia, Mod. Phys. Lett. A13, 2249 (1998).
- [8] Super-Kamiokande Collaboration, S. Fukuda *et al.*, Phys. Rev. Lett. 85, 3999 (2000).
- [9] R. Foot, Phys. Lett. B496, 169 (2000).
- [10] M. Shiozawa for the Super-Kamiokande Collaboration, talk at the Neutrino2002 conference, Munich, May 2002.
- [11] R. Foot, H. Lew and R. R. Volkas, Mod. Phys. Lett. A7, 2567 (1992); R. Foot, Mod. Phys. Lett. A9, 169 (1994); R. Foot and R. R. Volkas, Phys. Rev. D52, 6595 (1995).
- [12] Y. Itow *et al.*, hep-ex/0106019.